

# **RxHighlights**

May 2020

#### New drugs

Learn more

Drug name Manufacturer(s)	Therapeutic category	Indication(s)	Launch information
<b>Artesunate</b> <sup>†*</sup> Amivas	Antimalarial	Initial treatment of severe malaria in adult and pediatric patients	TBD
Darzalex Faspro <sup>™</sup> (daratumumab and hyaluronidase-fihj) Janssen	Humanized anti-CD38 monoclonal antibody	Treatment of adult patients with multiple myeloma: in combination with Velcade <sup>®</sup> (bortezomib), melphalan and prednisone in newly diagnosed patients who are ineligible for autologous stem cell transplant; in combination with Revlimid <sup>®</sup> (lenalidomide) and dexamethasone in newly diagnosed patients who are ineligible for autologous stem cell transplant and in patients with relapsed or refractory multiple myeloma who have received at least one prior therapy; in combination with bortezomib and dexamethasone in patients who have received at least one prior therapy, in patients who have received at least three prior lines of therapy including a proteasome inhibitor (PI) and an immunomodulatory agent	May 4, 2020
<b>Elyxyb</b> <sup>™</sup> (celecoxib) oral solution Dr. Reddy's	Nonsteroidal anti-inflammatory drug	Acute treatment of migraine with or without aura in adults	TBD

<b>Fensolvi<sup>®</sup></b> (leuprolide acetate) Tolmar Pharmaceuticals	Gonadotropin releasing hormone agonist	Treatment of pediatric patients 2 years of age and older with central precocious puberty	May 5, 2020
<b>Ferriprox</b> <sup>®</sup> (deferiprone) twice-a- day 1000 mg oral tablets Chiesi Global Rare Diseases	Iron chelator	Treatment of patients with transfusional iron overload due to thalassemia syndromes when current chelation therapy is inadequate	TBD
<b>Impeklo</b> <sup>™</sup> (clobetasol propionate) Mylan	Corticosteroid	Relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses in patients 18 years of age or older	TBD
<b>Kynmobi</b> <sup>™</sup> (apomorphine) Sunovion Pharmaceuticals	Non-ergoline dopamine agonist	Acute, intermittent treatment of "off" episodes in patients with Parkinson's disease	September 2020
Phexxi <sup>™</sup> (lactic acid/citric acid/potassium bitartrate)* Evofem	Spermicidal agent	Prevention of pregnancy in females of reproductive potential for use as an on-demand method of contraception	September 2020
Oriahnn <sup>™</sup> (elagolix/estradiol/norethindrone acetate; elagolix) AbbVie	Gonadotropin-releasing hormone receptor antagonist/estrogen/progestin	Management of heavy menstrual bleeding associated with uterine leiomyomas (fibroids) in premenopausal women	June 3, 2020
<b>Qinlock</b> <sup>™</sup> (ripretinib) <sup>†*</sup> Deciphera Pharmaceuticals	PDGFR-alpha kinase inhibitor	Treatment of adult patients with advanced gastrointestinal stromal tumor who have received prior treatment with 3 or more kinase inhibitors, including imatinib (eg, Gleevec <sup>®</sup> ).	May 19, 2020

<b>Retevmo</b> <sup>™</sup> (selpercatinib) <sup>†*</sup> Eli Lilly	RET inhibitor	Treatment of: adult patients with metastatic RET fusion- positive non-small cell lung cancer; adult and pediatric patients 12 years of age and older with advanced or metastatic RET-mutant medullary thyroid cancer who require systemic therapy; adult and pediatric patients 12 years of age and older with advanced or metastatic RET fusion-positive thyroid cancer who require systemic therapy and who are radioactive iodine-refractory (if radioactive iodine is appropriate)	May 11, 2020
<b>Tabrecta</b> ™ (capmatinib) <sup>†*</sup> Novartis	cMET inhibitor	Treatment of adult patients with metastatic non-small cell lung cancer whose tumors have a mutation that leads to mesenchymal-epithelial transition exon 14 skipping as detected by an FDA-approved test	May 10, 2020
VESIcare LS <sup>™</sup> (solifenacin succinate) oral suspension Astellas	Muscarinic antagonist	Treatment of neurogenic detrusor overactivity in pediatric patients aged 2 years and older	Late 2020
Zilxi <sup>™</sup> (minocycline) topical foam Menlo Therapeutics	Tetracyclines	Treatment of inflammatory lesions of rosacea in adults	Fourth quarter of 2020

\*New molecular entity †Orphan Drug TBD: To be determined

#### **New generics**

Learn more

Drug name Manufacturer(s)	Generic manufacturer(s)	Strength(s) & dosage form(s)	Therapeutic use	Launch information
<b>Samsca<sup>®</sup> (</b> tolvaptan) Otsuka	Alkem/Ascend <sup>†</sup>	30 mg tablets	Hypervolemic and euvolemic hyponatremia, including patients with heart failure and syndrome of inappropriate antidiuretic hormone	May 20, 2020

†A-rated generic manufacturer

#### New authorized brand alternatives

Drug name Manufacturer(s)	Generic manufacturer(s)	Strength(s) & dosage form(s)	Therapeutic use	Launch information
Aptensio XR <sup>™</sup> (methylphenidate) Rhodes Pharmaceuticals	Rhodes Pharmaceuticals	10 mg, 15 mg, 20 mg, 30 mg, 40 mg 50 mg, 60 mg extended-release capsules	Attention deficit hyperactivity disorder	May 5, 2020
<b>Samsca</b> <sup>®</sup> (tolvaptan) Otsuka	Apotex	30 mg tablets	Hypervolemic and euvolemic hyponatremia, including patients with heart failure and syndrome of inappropriate antidiuretic hormone	May 20, 2020

## Indications/label updates

Learn more

Drug name Manufacturer(s)	Туре	Description
<b>Alunbrig</b> <sup>®</sup> (brigatinib) Takeda	Expanded indication	Treatment of adult patients with anaplastic lymphoma kinase positive metastatic non-small cell lung cancer
<b>Brilinta<sup>®</sup></b> (ticagrelor) AstraZeneca	New indication	To reduce the risk of a first myocardial infarction or stroke in patients with coronary artery disease at high risk for such events
<b>Cyramza<sup>®</sup> (</b> ramucirumab) Eli Lilly	Expanded indication	In combination with erlotinib, for the first-line treatment of patients with metastatic non- small cell lung cancer whose tumors have epidermal growth factor receptor exon 19 deletions or exon 21 substitution mutations
<b>Dupixent<sup>®</sup></b> (dupilumab) Regeneron and Sanofi	Expanded indication	Treatment of patients aged 6 years and older with moderate-to-severe atopic dermatitis whose disease is not adequately controlled with topical prescription therapies or when those therapies are not advisable
<b>Farxiga</b> <sup>®</sup> (dapagliflozin) AstraZeneca	New indication	Reduce the risk of cardiovascular death and hospitalization for heart failure in adults with heart failure (New York Heart Association class II-IV) with reduced ejection fraction
<b>Lynparza<sup>®</sup></b> (olaparib) AstraZeneca and Merck	New indication	Treatment of adult patients with deleterious or suspected deleterious germline or somatic homologous recombination repair gene-mutated metastatic castration-resistant prostate cancer who have progressed following prior treatment with Xtandi <sup>®</sup> (enzalutamide) or abiraterone
<b>Opdivo</b> <sup>®</sup> (nivolumab) and <b>Yervoy</b> <sup>®</sup> (ipilimumab) Bristol Myers Squibb	New indications	As first-line treatment in combination with Yervoy in adult patients with metastatic non- small cell lung cancer expressing PD-L1 (≥ 1%) as determined by an FDA-approved test, with no EGFR or ALK genomic tumor aberrations Opdivo, in combination with Yervoy and 2 cycles of platinum-doublet chemotherapy, for the first-line treatment of adult patients with metastatic or recurrent non-small cell lung cancer, with no EGFR or ALK genomic tumor aberrations
<b>Pomalyst<sup>®</sup></b> (pomalidomide) Bristol Myers Squibb	New orphan indication	Treatment of adults with acquired immunodeficiency syndrome -related Kaposi sarcoma after failure of highly active antiretroviral therapy and Kaposi sarcoma in adult patients who are human immunodeficiency virus-negative

<b>Rubraca<sup>®</sup> (rucaparib)</b> Clovis Oncology	New indication	Treatment of adults with a deleterious BRCA mutation (germline and/or somatic) associated metastatic castration-resistant prostate cancer who have been treated with androgen receptor-directed therapy and a taxane-based chemotherapy
<b>Sirturo<sup>®</sup> (bedaquiline)</b> Janssen	Expanded indication	As part of combination therapy in the treatment of adult and pediatric patients (5 years and older and weighing at least 15 kg) with pulmonary multi-drug resistant tuberculosis (MDR-TB)
<b>Tecentriq</b> <sup>®</sup> (atezolizumab) Genentech	Expanded indication	As a single agent, for the first-line treatment of adult patients with metastatic non-small cell lung cancer whose tumors have high PD-L1 expression (PD-L1 stained $\geq$ 50% of tumor cells [TC $\geq$ 50%] or PD-L1 stained tumor-infiltrating immune cells [IC] covering $\geq$ 10% of the tumor area [IC $\geq$ 10%]), as determined by an FDA-approved test, with no EGFR or ALK genomic tumor aberrations
<b>Tecentriq</b> <sup>®</sup> (atezolizumab) and <b>Avastin</b> <sup>®</sup> (bevacizumab) Genentech	New indication	In combination with bevacizumab (eg, Avastin <sup>®</sup> ), is indicated for the treatment of patients with unresectable or metastatic hepatocellular carcinoma who have not received prior systemic therapy

# Drug safety news

Drug name Manufacturer(s)	Description
	The FDA announced that laboratory testing has revealed levels of N-Nitrosodimethylamine (NDMA) above the agency's acceptable intake limit in several lots of the extended-release formulation of metformin.
Metformin Extended-Release Products	The FDA is in contact with five firms to recommend they voluntarily recall their products. Company recall notices will be posted on FDA's website. There are additional manufacturers of the metformin ER formulation that supply a significant portion of the U.S. market, and their products are not being recalled. FDA testing has not shown NDMA in immediate release metformin products.

## Drug recalls/withdrawals/shortages/discontinuations

Learn more

Drug name Manufacturer(s)	Dosage form(s)	Туре	Description
Lactated Ringer's injection	1000 mL flexible container	Recall	ICU Medical announced a voluntary, consumer-level recall of one lot of Lactated Ringer's injection due to the presence of particulate matter identified as iron oxide. Lactated Ringer's injection is indicated for parenteral replacement of extracellular losses of fluid and electrolytes.
<b>NP Thyroid</b> <sup>®</sup> (thyroid tablets) Acella Pharmaceuticals	30 mg, 60 mg, and 90 mg tablets	Recall	Acella announced a consumer-level recall of thirteen lots of NP Thyroid because testing has found the recalled lots to be superpotent. The product may have up to 115.0% of the labeled amount of liothyronine. NP Thyroid is used to treat hypothyroidism.

## Key guideline/literature updates

Торіс	Reference
Eosinophilic Esophagitis Management - American Gastroenterological Association Institute and the American Academy of Allergy, Asthma & Immunology/American College of Allergy, Asthma & Immunology	<u>Gastroenterology</u> . May 2020
Ménière's Disease Executive Summary - American Academy of Otolaryngology - Head and Neck Surgery Foundation	<u>Otolaryngology – Head and Neck Surgery</u> . April 2020

# RxHighlights

Sickle Cell Disease: Prevention, Diagnosis, and Treatment of Cerebrovascular Disease in Children and Adults – American Society of Hematology	<u>Blood Advances</u> . April 2020
National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology:	<u>NCCN Clinical Practice Guidelines in Oncology: Anal Carcinoma</u> .
Anal Carcinoma - Version 2.2020	May 2020
National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology:	<u>NCCN Clinical Practice Guidelines in Oncology: Bladder Cancer</u> .
Bladder Cancer - Version 5.2020	May 2020
National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology:	<u>NCCN Clinical Practice Guidelines in Oncology: Breast Cancer</u> .
Breast Cancer - Version 4.2020	May 2020
National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology:	<u>NCCN Clinical Practice Guidelines in Oncology: Colon Cancer</u> .
Colon Cancer - Version 3.2020	May 2020
National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology: Cutaneous Melanoma - Version 3.2020	<u>NCCN Clinical Practice Guidelines in Oncology: Cutaneous</u> <u>Melanoma</u> . May 2020
National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology: Esophageal and Esophagogastric Junction Cancers - Version 2.2020	NCCN Clinical Practice Guidelines in Oncology: Esophageal and Esophagogastric Junction Cancers. May 2020
National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology:	<u>NCCN Clinical Practice Guidelines in Oncology: Gastric Cancer</u> .
Gastric Cancer - Version 2.2020	May 2020
National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology: Gestational Trophoblastic Neoplasia - Version 2.2020	<u>NCCN Clinical Practice Guidelines in Oncology: Gestational</u> <u>Trophoblastic Neoplasia</u> . May 2020

<u>NCCN Clinical Practice Guidelines in Oncology: Multiple Myeloma</u> . May 2020
<u>NCCN Clinical Practice Guidelines in Oncology: Myeloproliferative</u> <u>Neoplasms</u> . May 2020
<u>NCCN Clinical Practice Guidelines in Oncology: Non-Small Cell Lung</u> <u>Cancer</u> . May 2020
<u>NCCN Clinical Practice Guidelines in Oncology: Occult Primary</u> . May 2020
<u>NCCN Clinical Practice Guidelines in Oncology: Prostate Cancer</u> . May 2020
<u>NCCN Clinical Practice Guidelines in Oncology: Rectal Cancer</u> . May 2020
<u>NCCN Clinical Practice Guidelines in Oncology: Small Bowel</u> <u>Adenocarcinoma</u> . May 2020
<u>NCCN Clinical Practice Guidelines in Oncology: Soft Tissue Sarcoma</u> . May 2020
<u>NCCN Clinical Practice Guidelines in Oncology: Systemic</u> <u>Mastocytosis</u> . May 2020

National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology:	<u>NCCN Clinical Practice Guidelines in Oncology: Testicular Cancer</u> .
Testicular Cancer - Version 3.2020	May 2020
National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology:	<u>NCCN Clinical Practice Guidelines in Oncology: Uveal Melanoma</u> .
Uveal Melanoma - Version 1.2020	May 2020
National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology: Breast Cancer Risk Reduction - Version 1.2020	NCCN Clinical Practice Guidelines in Oncology: Breast Cancer Risk Reduction. May 2020
National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology: Cancer-Related Fatigue - Version 2.2020	<u>NCCN Clinical Practice Guidelines in Oncology: Cancer-Related</u> <u>Fatigue</u> . May 2020
National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology:	<u>NCCN Clinical Practice Guidelines in Oncology: Smoking Cessation</u> .
Smoking Cessation - Version 1.2020	May 2020

optum.com/optumrx

OptumRx specializes in the delivery, clinical management and affordability of prescription medications and consumer health products. We are an Optum<sup>®</sup> company — a leading provider of integrated health services. Learn more at **optum.com**.

All Optum<sup>®</sup> trademarks and logos are owned by Optum, Inc. All other brand or product names are trademarks or registered marks of their respective owners. This document contains information that is considered proprietary to OptumRx and should not be reproduced without the express written consent of OptumRx.

RxHighlights is published by the OptumRx Clinical Services Department.

© 2020 Optum, Inc. All rights reserved.