



OptumRx has partnered with CoverMyMeds to receive prior authorization requests, saving you time and often delivering real-time determinations.

Visit [go.covermymeds.com/OptumRx](http://go.covermymeds.com/OptumRx) to begin using this free service.

Please note: All information below is required to process this request.

Mon-Fri: 5am to 10pm Pacific / Sat: 6am to 3pm Pacific

## Repatha<sup>®</sup> Prior Authorization Request Form (Page 1 of 3)

DO NOT COPY FOR FUTURE USE. FORMS ARE UPDATED FREQUENTLY AND MAY BE BARCODED

Member Information (required)			Provider Information (required)		
Member Name:			Provider Name:		
Insurance ID#:			NPI#:		Specialty:
Date of Birth:			Office Phone:		
Street Address:			Office Fax:		
City:	State:	Zip:	Office Street Address:		
Phone:			City:	State:	Zip:
Medication Information (required)					
Medication Name:			Strength:		Dosage Form:
<input type="checkbox"/> Check if requesting <b>brand</b>			Directions for Use:		
<input type="checkbox"/> Check if request is for <b>continuation of therapy</b>					
Clinical Information (required)					
<b>Select the diagnosis below:</b>					
<input type="checkbox"/> Homozygous familial hypercholesterolemia (HoFH)					
<input type="checkbox"/> Prevention of cardiovascular events					
<input type="checkbox"/> Primary hyperlipidemia					
<input type="checkbox"/> Atherosclerotic cardiovascular disease (ASCVD)					
<input type="checkbox"/> Heterozygous familial hypercholesterolemia (HeFH)					
<input type="checkbox"/> Other diagnosis: _____ ICD-10 Code(s): _____					
<b>Medication History:</b>					
Select if the patient has had history of failure, contraindication, or intolerance to the following:					
<input type="checkbox"/> Atorvastatin calcium		<input type="checkbox"/> Juxtapid		<input type="checkbox"/> Pravastatin sodium	
<input type="checkbox"/> Ezetimibe		<input type="checkbox"/> Lovastatin		<input type="checkbox"/> Rosuvastatin calcium	
<input type="checkbox"/> Ezetimibe/simvastatin		<input type="checkbox"/> Praluent		<input type="checkbox"/> Simvastatin	
<b>Clinical Information:</b>					
Select if the requested medication is prescribed by or in consultation with one of the following specialists:					
<input type="checkbox"/> Cardiologist					
<input type="checkbox"/> Endocrinologist					
<input type="checkbox"/> Lipid specialist					
Will Repatha be used in combination with another proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor? <input type="checkbox"/> Yes <input type="checkbox"/> No					
Does the prescriber attest that the information provided is true and accurate to the best of their knowledge and they understand that UnitedHealthcare may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided? <input type="checkbox"/> Yes <input type="checkbox"/> No					
<b>For homozygous familial hypercholesterolemia, also answer the following:</b>					
Select if the patient has homozygous familial hypercholesterolemia (HoFH) as confirmed by the following:*					
<input type="checkbox"/> Genetic confirmation of 2 mutations in the LDL receptor, ApoB, PCSK9, or LDL receptor adaptor protein 1 (i.e., LDLRAP1 or ARH)					
<input type="checkbox"/> Untreated LDL-C level greater than 500mg/dL					
<input type="checkbox"/> Treated LDL-C level greater than 300mg/dL					
<input type="checkbox"/> Xanthoma before 10 years of age					
<input type="checkbox"/> Evidence of heterozygous familial hypercholesterolemia in both parents					
*Please note: Chart documentation of the above is required to be submitted along with this fax.					
Is the patient receiving other lipid lowering therapy (e.g., statin, ezetimibe)? <input type="checkbox"/> Yes <input type="checkbox"/> No					
Will Repatha be used in combination with Juxtapid (Iomitapide)? <input type="checkbox"/> Yes <input type="checkbox"/> No					
Will Repatha be used in combination with Kynamro (mipomersen)? <input type="checkbox"/> Yes <input type="checkbox"/> No					

This document and others if attached contain information that is privileged, confidential and/or may contain protected health information (PHI). The Provider named above is required to safeguard PHI by applicable law. The information in this document is for the sole use of OptumRx. Proper consent to disclose PHI between these parties has been obtained. If you received this document by mistake, please know that sharing, copying, distributing or using information in this document is against the law. **If you are not the intended recipient, please notify the sender immediately.**

Office use only: Repatha\_CMS\_2019Feb-W

## Repatha<sup>®</sup> Prior Authorization Request Form (Page 2 of 3)

DO NOT COPY FOR FUTURE USE. FORMS ARE UPDATED FREQUENTLY AND MAY BE BARCODED

### For prevention of cardiovascular events and primary hyperlipidemia, also answer the following:

Select the indication for which this medication is being prescribed:

- Heterozygous familial hypercholesterolemia (HeFH) as confirmed by the following:
  - Untreated/pre-treatment LDL-cholesterol (LDL-C) greater than 190 mg/dL as an adult
  - Family history of myocardial infarction in first-degree relative less than 60 years of age
  - Family history of myocardial infarction in second-degree relative less than 50 years of age
  - Family history of LDL-C greater than 190 mg/dL in first- or second-degree relative
  - Family history of familial hypercholesterolemia in first- or second-degree relative
  - Family history of tendinous xanthomas and/or arcus cornealis in first- or second-degree relative
  - Presence of tendinous xanthomas in the patient\*
  - Arcus cornealis before age 45\*
  - Functional mutation in the LDL receptor, ApoB, or PCSK9 gene\*

\*Please note: Chart documentation of the above is required to be submitted along with this fax.

- Atherosclerotic cardiovascular disease (ASCVD) as confirmed by the following:
  - Acute coronary syndromes
  - Coronary or other arterial revascularization
  - History of myocardial infarction
  - Stroke
  - Stable or unstable angina
  - Transient ischemic attack
  - Peripheral arterial disease presumed to be of atherosclerotic origin
- Prevention of cardiovascular events

Select if the patient has one of the following LDL-C values while on maximally tolerated lipid-lowering regimen **within the last 120 days**:

- LDL-C between 70 mg/dL and 99 mg/dL **with** ASCVD
- LDL-C greater than or equal to 100 mg/dL **with** ASCVD
- LDL-C between 100 mg/dL and 129 mg/dL **without** ASCVD
- LDL-C greater than or equal to 130 mg/dL **without** ASCVD

Has the patient been receiving at least 12 consecutive weeks of one **high-intensity** statin therapy [i.e., atorvastatin 40-80 mg, Crestor (rosuvastatin) 20-40 mg] at a maximally tolerated dose?  Yes  No

Will the patient continue to receive a **high-intensity** statin therapy?  Yes  No

Select if the patient is unable to tolerate **high-intensity** statin therapy as evidenced by one of the following intolerable and persistent (i.e., more than 2 weeks) symptoms:

- Myalgia (muscle symptoms without CK elevations)
- Myositis (muscle symptoms with CK elevations less than 10 times upper limit of normal [ULN])

Has the patient been receiving at least 12 consecutive weeks of one **moderate-intensity or low-intensity** statin therapy [i.e., atorvastatin 10-20 mg, Crestor (rosuvastatin) 5-10 mg, simvastatin 10-40 mg, pravastatin 10-80 mg, lovastatin 20-40 mg, Lescol XL (fluvastatin XL) 80 mg, fluvastatin 20-40 mg, or Livalo (pitavastatin) 1-4 mg] at a maximally tolerated dose?  Yes  No

Will the patient continue to receive a **moderate-intensity or low-intensity** statin therapy?  Yes  No

Select if the patient is unable to tolerate **moderate-intensity or low-intensity** statin therapy as evidenced by one of the following intolerable and persistent (i.e., more than 2 weeks) symptoms:

- Myalgia (muscle symptoms without CK elevations)
- Myositis (muscle symptoms with CK elevations less than 10 times upper limit of normal [ULN])

Does the patient have a labeled contraindication to all statins as documented in medical records?\*  Yes  No

\*Please note: Chart documentation of the above is required to be submitted along with this fax.

Has the patient experienced rhabdomyolysis or muscle symptoms with statin treatment with CK elevations greater than 10 times ULN on one statin therapy?  Yes  No

Has the patient been receiving at least 12 consecutive weeks of ezetimibe (Zetia) therapy as adjunct to maximally tolerated statin therapy?  Yes  No

If "no to the above question, does the patient have a history of contraindication or intolerance to ezetimibe (Zetia)?  Yes  No



## Repatha® Prior Authorization Request Form (Page 3 of 3)

DO NOT COPY FOR FUTURE USE. FORMS ARE UPDATED FREQUENTLY AND MAY BE BARCODED

### Reauthorization:

If this is a reauthorization request, answer the following questions:

Has the patient experienced LDL-C reduction while on Repatha therapy as documented in medical records (e.g., chart notes, laboratory values)?\*  Yes  No

\*Please note: Chart documentation of the above is required to be submitted along with this fax.

Select if the requested medication is prescribed by or in consultation with one of the following specialists:

- Cardiologist       Endocrinologist       Lipid specialist

Will Repatha be used in combination with another proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor?  Yes  No

Does the prescriber attest that the information provided is true and accurate to the best of their knowledge and they understand that UnitedHealthcare may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided?  Yes  No

For primary hyperlipidemia, also answer the following questions:

Does the patient continue to receive a statin at the maximally tolerated dose (unless patient has documented inability to take statins)?  Yes  No

For homozygous familial hypercholesterolemia, also answer the following questions:

Does the patient continue to receive other lipid-lowering therapy (e.g., statin, ezetimibe)?  Yes  No

Will Repatha be used in combination with Juxtapid (lomitapide)?  Yes  No

Will Repatha be used in combination with Kynamro (mipomersen)?  Yes  No

### Quantity Limit:

What is the quantity requested per MONTH? \_\_\_\_\_

What is the reason for exceeding the plan limitations?

- Titration or loading-dose purposes
- Patient is on a dose-alternating schedule (e.g., one tablet in the morning and two tablets at night, one to two tablets at bedtime)
- Requested strength/dose is not commercially available
- There is a medically necessary justification why the patient cannot use a higher commercially available strength to achieve the same dosage and remain within the same dosing frequency. **Please specify:** \_\_\_\_\_
- Other: \_\_\_\_\_

Are there any other comments, diagnoses, symptoms, medications tried or failed, and/or any other information the physician feels is important to this review?

### Please note:

This request may be denied unless all required information is received.

If the patient is not able to meet the above standard prior authorization requirements, please call 1-800-711-4555.

For urgent or expedited requests please call 1-800-711-4555.

This form may be used for non-urgent requests and faxed to 1-800-527-0531.